
REAC

S4414 Succession Administration



FFICE OF EMERGENCY AND REMEDIAL RESPONSE

NLI 002 0429

FINAL REPORT

TCLP SCREENING

NATIONAL LEAD INDUSTRIES SITE PEDRICKTOWN, NJ

FEBRUARY, 1993



PREPARED BY:

Mark D. Sprenger, Ph.D. Environmental Response Team

AND

Paul Bovitz ERT/REAC

Environmental Response Branch Emergency Response Division Office of Emergency & Remedial Response

TABLE OF CONTENTS

LIST OF TABLES

LIST OF FIGURES

EXECUTIVE SUMMARY

1.0 INTRODUCTION

- 1.1 Objective of this Study
- 1.2 Site Background

2.0 METHODOLOGY

- 2.1 Field Sampling Design
- 2.2 XRF Screening
- 2.3 Soil and Sediment Sampling
- 2.4 Analyses
- 2.5 Data Analysis and Interpretation

3.0 RESULTS

- 3.1 XRF Screening
- 3.2 Metals in Soil and Sediment Measured by AA
 - 3.2.1 Lead
 - 3.2.2 Other Metals
- 3.3 TCLP Metals Extracts from Soil and Sediment
 - 3.3.1 TCLP Results for Pb
 - 3.3.2 TCLP Results for As, Ba, Cr and Cd
- 3.4 TOC of Soil and Sediment
- 3.5 Grain Size of Soil and Sediment

4.0 DISCUSSION

- 4.1 TCLP Results of Lead-Contaminated Soil and Sediment
- 4.2 TCLP Results for Other Metals in Soil and Sediment

5.0 CONCLUSIONS

REFERENCES

APPENDICES

Appendix A - REAC Analytical Report

Appendix B - U.S. EPA ERT/TAT XRF Results

Appendix C - Statistical Output

LIST OF TABLES

TABLE	
1	Soil and Sediment Sampling Locations for TCLP Analysis
2	Lead Concentrations in Soil and Sediment as Measured by XRF and AA
3	Lead Concentrations in Soil and Sediment, and Corresponding TCLP Extractions
4	Arsenic and Barium Concentrations in Soil and Sediment as Measured by AA, and Corresponding Concentrations in TCLP Extracts
5	Cadmium and Chromium Concentrations as Measured by AA, and Corresponding Concentrations in TCLP Extracts
6	Maximum TCLP Concentrations of Metals as Compared to RCRA Regulatory Limits
7	Results of TOC Analysis of Soil and Sediment Samples
8	Results of Grain Size Analysis of Soil and Sediment

LIST OF FIGURES

FIGURE	
1	Site Location Map
2	XRF Screening and TCLP Soil and Sediment Sampling Locations

1.0 INTRODUCTION

1.1 Objective of this Study

The objective of this project was to evaluate representative contaminanted soils and sediment at the National Lead (NL) site, with respect to compliance with Resource Conservation and Recovery Act (RCRA) criteria for classification as a hazardous material through TCLP procedures.

1.2 Site Background

The NL Industries site is located in Pedricktown, Salem County, New Jersey (Figure 1). The site is an abandoned secondary lead smelting facility located approximately 1.5 miles east of the Delaware River. The facility is situated on 44 acres and was operated from 1972 to 1984. Lead batteries and other lead materials were handled by this facility. Site contaminants include lead (Pb), cadmium (Cd), and chromium (Cr).

The area of investigation roughly included the area between two streams running in a northerly direction along both sides of the former facility, and areas mapped as forested wetland which are associated with these drainages⁽¹⁾. Concentrations of Pb in surface soil range from 35 milligrams per kilogram (mg/kg) to above 20,200 mg/kg⁽¹⁾.

2.0 METHODOLOGY

2.1 Field Sampling Design

Twenty-seven sample locations in the area immediately surrounding the NL Industries site were screened for Pb using X-Ray Fluorescence Spectrometry (XRF) (Figure 2). Based upon these results, 18 of these locations were actually sampled. These consisted of 14 soil samples and four sediment samples. The samples were collected across a range of concentrations in the West Stream and within four general habitats, each with different soil types. Four samples were collected from the West Stream (sediment/muck), four from the upland forest (sandy loam), five from the forested wetland (sandy loam with high organic content), and five from the facility area (loamy sand).

2.2 XRF Screening

Upland and facility soil samples were screened using XRF in situ. Previously sampled areas were included in order to minimize time and effort spent in locating desired

soil concentrations. If, upon screening, a sample was within the desired concentration range, it was collected with a decontaminated stainless steel trowel and homogenized before placement into glass jars.

Wetland and stream samples were collected using a decontaminated stainless steel bucket auger or trowel, homogenized in an aluminum pan, and apportioned into labeled glass jars. A 2-ounce aliquot from each jar was dried and screened using XRF. Samples which were selected were retained in glass jars on ice. Otherwise a sample was returned to its original location.

2.3 Soil and Sediment Sampling

At each location selected for analysis, three jars were filled with the homogenized soil. These included one 32-ounce jar for grain size analysis and one 16-ounce jar for Atomic Absorption (AA) analysis of metals and Toxicity Characteristic Leachate Procedure (TCLP) Metals. A 2-ounce jar for Total Organic Carbon (TOC) analysis was also collected. At locations with wet soils, an additional 2-ounce jar was filled for XRF screening according to the procedure above (section 2.2).

Field sampling activities were conducted according to the following U.S. EPA ERT/REAC Standard Operating Procedures (SOPs): SOP #1713, Spectrace 9000 Field Portable X-Ray Fluorescence Operating Procedures; SOP #2012, Soil Sampling; SOP #2016, Sediment Sampling; SOP #2125, Trowels and Scoops, and SOP #2127, Hand Bucket Augers.

All sampling equipment was decontaminated between locations. Decontamination procedures followed the sequence below:

- Soil and other material were physically removed.
- Equipment was washed with soap and potable water.
- Equipment was rinsed with distilled water.
- Equipment was sprayed with 10 percent nitric acid.
- Equipment was rinsed again with distilled water.
- Equipment was allowed to air dry.

Sample handling and documentation followed U.S. EPA ERT/REAC SOPs #2002, Sample Documentation; SOP #4004, Chain of Custody, and SOP #2003, Sample Storage, Preservation and Handling.

2.4 Analyses

All analyses were conducted by a subcontracting laboratory (ATEC Consultants, Inc., Indianapolis, IN). Eight metals on the TCLP list, including Cd, Cr, Pb, arsenic (As), barium (Ba), mercury (Hg), selenium (Se), and silver (Ag), were analyzed using AA (procedures in Appendix A). After an aliquot was taken for AA analysis, the remainder of the sample was submitted for TCLP analyses. Grain size and TOC analyses were also conducted.

2.5 Data Analyses and Interpretation

In order to determine if metals were collocated, the concentrations of metals which were found above their respective detection limits were analyzed using correlation analysis.

All metals found above their respective method detection limits were analyzed to determine the percentage of metal leached during the TCLP analysis using the following procedure.

- Analytical results were converted to a dry weight basis, so that comparisons were not influenced by the moisture content of the sample. Since moisture content could be expected to change drastically with field conditions, e.g., a wetland may not always be wet, conversion to dry weight allows a common basis for comparison of soil/sediment leaching ability between sample locations.
- The amount of metal (in mg) present in the 100-g sample used for the TCLP extraction procedure was determined by multiplying the dry weight concentration (mg/kg) by 0.1 kg, since there are 1000 g in a kg.
- The amount of metal (in mg) extracted from the 100g sample was calculated by taking the measured concentration in mg/L and multiplying it by 2 L, the amount of solution used to extract the metal.
- This quantity was divided by the original amount of metal present in the soil (mg) to determine the percent of the original metal lost to leaching during the procedure.

3.0 RESULTS

3.1 XRF Screening

Results of XRF screening for Pb revealed surface concentrations ranging from 230 mg/kg to 19,180 mg/kg (Table 1 and Appendix B). Concentrations in the upland forest located off site and east of the facility (Points E12-E29) ranged from 370 to 1230 mg/kg. A high concentration of 3220 mg/kg was recorded at sample point EE1A, located within the "hotspot" in the forested wetland southeast of the facility (Figure 2).

Sample points screened within the immediate vicinity of the facility (P1-P5) ranged in concentration from 230 to 10,000 mg/kg Pb. Points selected in the forested wetland located west of the facility (points W2-W20) had concentrations ranging from 730 to 19,180 mg/kg (Table 1 and Figure 2). Sediment samples collected from the West Stream (S2-S13) ranged in concentration from 270 to 17,930 mg/kg.

XRF concentrations determined in the field (in situ in the case of soils) were notably higher than those measured by AA in the samples actually collected (Table 2). However, a regression of AA versus XRF results indicated the two data sets were positively correlated, and a regression analysis indicated a significant linear relationship between AA and XRF results (p<0.01). Nevertheless, there was significant variability evident $(r^2=0.38, Appendix C)$.

3.2 Metals in Soil and Sediment Measured by AA

3.2.1 Lead

At all locations, Pb had the highest concentration of any metal analyzed by AA, with concentrations ranging from 96 to 16,000 mg/kg (wet weight). Wet weight Pb concentrations were correlated with those of As, Ba and Cd (r=0.63, r=0.66, and r=0.75, respectively), indicating these metals are collocated to some degree.

3.2.2 Other Metals

Of the remaining metals on the TCLP list, only Ba and Cr were found in all samples. Arsenic was found in all but one sample, while Cd was detected in approximately half the samples. Mercury, Se, and Ag were undetected at their respective detection limits in all samples (Appendix A, Table 1.1).

Barium was found in concentrations ranging from 6.8 to 87 mg/kg wet weight (Appendix A). The wet weight Ba concentration of 87 mg/kg at location W18 was substantially higher than that of the other sample locations. In addition to being correlated with Pb, wet weight Ba concentrations were significantly correlated with those of As (r=0.77, Appendix C).

Chromium was detected at values ranging from 4.8 to 38 mg/kg (wet weight). At 38 mg/kg wet weight, the concentration of Cr was substantially higher at location EE1A, the hotspot in the forested wetland southeast of the site, than at other locations. Concentrations of Cr were not correlated with any other metal.

Cadmium concentrations ranged from undetected to a maximum of 23 mg/kg wet weight. The latter value was recorded at location W18. Cadmium was significantly correlated only with Pb.

3.3 TCLP Metals Extracts from Soil and Sediment

3.3.1 TCLP Results for Pb

Lead was the only metal tested for which TCLP concentrations exceeded regulatory criteria under RCRA⁽³⁾. For Pb, samples from which the TCLP extract concentrations at or above 5 mg/L are considered hazardous materials under RCRA. This criteria value was exceeded in TCLP extracts from sample locations P3, P5, W18, W20, S13, and S11 (Table 3).

Results of TCLP analysis of Pb-contaminated soil and sediment showed the degree of leaching to vary with the original Pb concentration. The amount of Pb present in the TCLP extract was correlated with wet weight Pb concentrations in soil and sediment (r=0.84), and this relationship was highly significant (p<0.001), Appendix C). Conversion to dry weight concentrations lowered the correlation coefficient to 0.69, but the relationship was still significant (p<0.001).

Review of the data in Table 3, with the exception of location P2, indicates that the soil or sediment samples with less than 1000 mg/kg Pb had TCLP values of less than 1 mg/L Pb. A regression analysis using soil/sediment Pb concentration (dry weight) as the independent variable and Pb concentration in TCLP extract as the dependent variable showed the relationship to be exponential $(r^2 = 0.84, p < 0.01)$.

The percentage of Pb from the original sample which leached during the TCLP analyses ranged from <1 percent to approximately 54 percent. Samples from locations P2 and P5 had a significantly higher percentage of Pb leached, with approximately 54 and 53 percent, respectively. Yet, while location P5 had the second highest Pb concentration of any soil sample (17,391 mg/kg, dry weight), the soil collected from location P2 had a Pb concentration measured at only 107 mg/kg dry weight.

The amount of Pb leached from the soil varied between areas of the site investigated (Tables 1 and 3). Samples collected from the woodland on the eastern side of the facility (E12-E29) and from the "hotspot" in the woods southeast of the facility (EE1A) ranged in Pb contamination from 325 to 927 mg/kg (dry weight). However, the percent of Pb leached in the TCLP analyses was under 3 percent in all six samples.

Samples collected in the immediate vicinity of the facility (P1-P5) ranged in Pb contamination from 107 to 17391 mg/kg dry weight and exhibited TCLP values ranging from approximately 5 to 54 percent. Samples collected from the forested wetland on the western side of the facility had a similar Pb contamination range (185 mg/kg to 18987 mg/kg dry weight), but the percentage leached ranged only from <1 to 12.64 percent. Finally, results of TCLP analyses of samples collected from the West Stream (S2-S13) indicate that while Pb contamination ranged from 368 to 13611 mg/kg dry weight, the percentage of Pb leached during the TCLP analyses ranged from only 1.57 to 6.32 percent.

3.3.2 TCLP Results for As, Ba, Cr, and Cd.

Arsenic was detected in the leachate of only two samples: S11 and S13 (Table 4). The percent of the original amount present which leached varied from ND to 12.96 percent. The maximum concentration of As leached was 0.9 mg/L.

Barium was detected in the leachate of all but four samples (E16-E29, and EE1A); these all had soil concentrations of less than 12 mg/kg (Table 4). The highest amount of Ba leached (1 mg/L) was from sample W2. The percentage of Ba leached from samples ranged from undetected to nearly 34 percent (sample location W2). The percentage of Ba leached from the sample was highly variable and did not appear to be related to the concentration in the soil or sediment. For instance, the highest

proportion leached (33.75 percent) was from sample location W2, where the soil concentration was 32 mg/kg. At locations W18 and W20 the soil concentrations were notably higher (87 and 47 mg/kg, respectively), yet the amount leached was substantially lower (5.99 and 5.17 percent, respectively).

Chromium was recorded in the leachate from only one sample (W18), at a concentration of 0.1 mg/L (Table 5). This represented 5.83 of the Cr in the soil sample. Cr was not detected in the leachate from two other locations (W20 and EE1A) where soil concentrations were notably higher.

Cadmium was found in the leachate of six of the 18 samples submitted for analysis. The maximum Cd level recorded was 0.4 mg/L, in sample number W18 (Table 5). The percentage of Cd released into solution did not appear to be directly related to soil or sediment concentrations; three samples with varying concentrations (P4, P5, and W18) had a similar percentage of Cd leached (from 25-27 percent, Appendix C). Maximum TCLP concentrations for all metals and corresponding maximum permissable levels under RCRA are presented in Table 6.

3.4 TOC of Soil and Sediment

TOC levels were below 30,000 mg/kg, the upper limit of the analysis, at locations P1 through P5 (Table 7). At these locations, TOC concentrations varied from 2,650 to 7,675 mg/kg. Exact concentrations at levels above 30,000 mg/kg could not be calculated by the subcontracted laboratory due to the high amount of organic material present in the soil and sediment at those locations. However, analyses conducted in prior field a investigation by U.S. EPA/ERT and REAC(2) using alternative analytical method found levels ranging from 66,000 to 160,000 mg/kg in West Stream samples. In the same study, TOC values ranged from 13,000 to 68,000 mg/kg in soils collected in the woodland area east of the facility, and from 41,000 to 200,000 mg/kg in soils from the forested wetland west of the facility.

3.5 Grain Size of Soil and Sediment

Analysis of particle size distribution showed that the 18 soil and sediment samples collected consisted mostly of sand. Sand content ranged from 42.4 to 87.2 percent, with the majority of samples consisting of more than 50 percent sand (Table 8). Most of the sand fraction consisted of medium and fine sand. The proportion of

sand in samples from the eastern woodland (locations E12-E29) was consistently between 80 and 86 percent. Soil samples from the forested wetland west of the site had lower amounts of sand, which ranged between 40 and 80 percent. Samples collected within the immediate vicinity of the facility (P1-P4) had variable amounts of sand, ranging from about 60 to 87 percent. The sample collected from the woodland area southeast of the facility consisted of approximately 53 percent sand. Sediment samples collected from the West Stream (S2-S13) ranged from approximately 70 to 85 percent sand.

The silt fraction was highest (41.3 percent) in the soil sample collected from the southeastern woodland (EE1A). Both silt and clay percentages were highly variable in samples collected throughout the site; silt values ranged from approximately 10 to 41 percent, and clay values ranged from approximately 1 to 23 percent. Differences in silt/clay fractions between areas generally reflected the differences in sand content described above.

4.0 DISCUSSION

4.1 TCLP Results of Lead-Contaminated Soil and Sediment

Lead concentrations in leachate from the soil and sediment exceeded RCRA regulatory criteria at six sample locations: P3, P5, W18, W20, S11, and S13. These locations were scattered throughout the site; S11 and S13 were taken from the West Stream, W18 came from the forested wetland west of the facility, and P3 and P5 were from the facility area itself. These samples had the four highest Pb concentrations of the 18 samples analyzed.

The above results are consistent with a general trend in this investigation; as the initial Pb concentration of soil and sediment samples increased, a logarithmic increase was observed in the amount of Pb leached into solution. This was true despite some obvious differences in the grain size and TOC characteristics of the soils which were measured.

Although there is a definite trend toward increased leaching with contaminant concentration (see Section 3.3.1 above), comparison of individual samples with similar concentrations showed that they each leached a different proportion of the original Pb present. For example, soil from W18 had a concentration of 18,987 mg/kg Pb and leached 12.64 percent of the total Pb present, while soil from P5 had a concentration of 17,391 mg/kg Pb and leached 52.9 percent of its Pb. Sample locations within the immediate vicinity of the facility, P2, P3, and P5, exhibited the highest percentage of Pb

leached, however, only the second two locations exceeded RCRA criteria.

The proportion of Pb leached does not appear to be related to the grain size distribution of the soil. Similarly, TOC showed no apparent relationship with the percentage of Pb leached, although comparisons were limited by the censored nature of the data.

4.2 TCLP Results for Other Metals in Soil and Sediment

Arsenic was present in the leachate of only two sediment samples (S11 and S13), both of which were collected from the West Stream. Only about 13 percent of the original was leached from these samples, with less than 1 mg/L was detected in solution. A maximum concentration of 139 mg/kg As (wet weight) was found in the sediment.

Barium was detected in the leachate of all but four samples (E16, E17, E29, and EE1A). However, the maximum Ba concentration detected in leachate (1 mg/L) was well below the RCRA standard of 100 mg/L.

Chromium was found in the leachate from only one sample, location W18, but was well below the RCRA limit. No Cr was detected in the leachate of the sample with the highest initial Cr concentration (93 mg/kg), which was from location EE1A.

Cadmium was present in less than half of the soil and sediment samples, and it was detected in the leachate of only six samples. With a maximum leachate concentration of 0.5 mg/L, it is well within RCRA regulatory limits.

The remaining metals, silver (Ag), mercury (Hg), and selenium (Se), were undetected in the soil, sediment, and leachate.

5.0 CONCLUSIONS

- (1) Lead concentrations in leachate, as measured by the TCLP procedure, exceeded RCRA regulatory criteria in sediment and soil from six of 18 sample locations (P3, P5, W18, W20, S11, and S13). These were located near the facility, in the forested wetland west of the facility, and in the West Stream (Figure 2).
- (2) Lead concentrations in leachate increased logarithmically with the original sample concentration in the soil and sediment.
- (3) Lead concentrations in leachate showed little relationship with grain size distribution or TOC content of soil and sediment samples.

(4) Remaining metals investigated (Ag, As, Ba, Cd, Cr, Hg, Se) did not exceed RCRA criteria in leachate from any sample location.

REFERENCES

- O'Brien and Gere. 1990. Remedial Investigation for National Smelting of New Jersey, Inc./NL Industries, Inc. Site. Volumes I-V. Unpubl. Manuscript.
- U.S. EPA ERT/REAC, 1992. Ecological Assessment of the National Lead Industries Site, Pedricktown, N.J. (Draft Document, Do Not Cite or Quote).
- (3) Federal Register, Vol. 55, no. 61. Rules and Regulations. 1990.

Tables

TABLE 1. Soil and Sediment Sampling Locations for TCLP Analysis NL Industries Site December, 1992

Habitat/Location	Sample Location	XRF Measured Pb Concentration (mg/kg) ⁽¹⁾	Soil Type ⁽²⁾
Forested Upland	E12	370	Sandy loam; little organic.
(East)	E29	30	
	E16	710	
	E17	1230	
Forested Wetland (Southeast)	EE1A	3220	Loam with high organic.
Facility Area	P2	230	Fine Sand.
	. P1	410	
	P4	3780	
	Р3	7040	
	P5	10000	
Forested Wetland	W2	730	Highly organic.
(West)	W12	1400	
	W18	10480	
	W 20	19180	
West Stream Sediment	S2	270	Muck.
	S4	910	
	S11	5670	
	S13	17930	

⁽¹⁾ All values rounded to the nearest 10¹ place. (2) General field observations.

TABLE 2. Lead Concentrations in Soil and Sediment as Measured by XRF and AA NL Industries Site
December, 1992

Sample Location	XRF Results (mg/kg)	AA Results (mg/kg wet weight)
E12	370	380
E29	530	270
E16	710	440
E17	1225	420
EE1A	· 3220	380
P1	410	170
P2	230	96
Р3	7040	3600
P4 -	3780	1000
P5	10000	16000
W2	730	100
W12	1400	780
W18	10480	15000
W20	19180	5100
S2	270	140
S4	910	730
S11	5670	2200
S13	17930	4900

TABLE 3. Lead Concentrations in Soil and Sediment, and Corresponding TCLP Extractions
NL Industries Site
December, 1992

Sample Location	Soil/Sediment Lead Concentration (mg/kg dry weight)	Pb Concentration in TCLP Extract (mg/L)	Percent Pb Extracted
E12	487	0.72	2.96
E29	325	0.42	2.58
E16	543	0.57	2.10
E17	494	0.55	2.23
EE1A	927	0.45	0.97
P2	107	2.90	54.38
P1 -	191	0.48	5.03
P4	1111	3.30	5.94
Р3	3956	43	21.74
P5	17391	460	52.90
W2	185	0.24	2.60
W12	2294	0.90	0.78
W18	18987	120	12.64
W20	11333	18	3.18
S2	368	0.96	5.21
S4	1780	1.4	1.57
S11	5789	11	3.80
S13	13611	43	6.32

Boldface locations represent those which had leachate values in exceeding RCRA criteria. MDL for the TCLP analysis was $0.05\ mg/L$.

TABLE 4. Arsenic and Barium Concentrations in Soil and Sediment as Measured by AA, and Corresponding Concentrations in TCLP Extracts NL Industries, Inc. Site December, 1992

Sample Location	Arsenic Concentr- ation (mg/kg dry wt)	Arsenic TCLP Extract (mg/L)	Percent Arsenic Extracted	Barium Concentr- ation (mg/kg dry wt)	Barium TCLP Extract (mg/L)	Percent Barium Extracted
E12	6.0	ND ⁽²⁾	•	23.0	0.12	10.4
E29	6.0	ND	*	11.0	ND ⁽³⁾	•
E16	8.0	ND	•	11.0	ND	*
E17	6.0	ND	*	8.0	ND	
EE1A	13.0	ND	•	34.0	ND	*
P2	3.0	ND	•	14.0	0.23	31.85
P1	6.0	ND	*	26.0	0.14	10.83
P4	11.0	ND	*	13.0	0.1	14.25
Р3	21.0	ND	*	30.0	0.07	4.65
P5	15.0	ND	*	33.0	0.1	6.13
W2	13.0	ND	*	59.0	1.0	33.8
W12	56.0	ND	•	76.0	0.11	2.88
W18	66.0	ND	*	110.0	0.33	5.99
W20	124.0	ND	*	104.0	0.3	5.17
S2	ND ⁽¹⁾	ND	*	26.0	0.1	4.18
S4	10.0	ND	•	59.0	0.21	7.17
S11	79.0	0.5	12.67	39.0	0.16	8.11
S13	139.0	0.9	12.96	86.0	0.7	15.79

⁽¹⁾ Arsenic method detection limit = 2.0 mg/kg
(2) Arsenic TCLP method detection limit = 0.50 mg/L
(3) Barium TCLP method detection limit = 0.50 mg/L
*Indicates percentage was not calculated due since compound was undetected in leachate.

TABLE 5. Cadmium and Chromium Concentrations as Measured by AA, and Corresponding Concentrations in TCLP Extracts
NL Industries Site December, 1992

Sample Location	Cadmium Concentration (mg/kg dry wt)	Cadmium TCLP Extract (mg/L)	Chromium Concentration (mg/kg dry wt)	Chromium TCLP Extract (mg/L)
E12	ND ⁽¹⁾	ND ⁽²⁾	11.0	ND ⁽³⁾
E29	ND	ND	9.0	ND
E16	ND	ND	11.0	ND.
E17	ND	ND	9.0	ND
EE1A	ND	ND	93.0	ND
P2	ND	ND	18.0	ND
P1	ND	ND	11.0	ND
P4	4.0	0.5	7.0	ND
Р3	2.0	ND	11.0	ND
P5	22.0	0.3	15.0	ND
W2	ND	ND	9.0	ND
W12	ND	ND	21.0	ND
W18	29.0	0.4	33.0	0.10
W20	10.0	0.03	44.0	ND
S2	ND	ND	13.0	ND
S4	5.1	ND	19.0	ND
S11	26.0	0.07	13.0	ND
S13	19.0	0.07	19.0	ND

 $^{^{(1)}} Cadmium$ method detection limit = 2 mg/kg $^{(2)} Cadmium$ TCLP method detection limit = 0.01 mg/L $^{(3)} Chromium$ TCLP method detection limit = 0.05 mg/L

TABLE 6. Maximum TCLP Concentrations of Metals as Compared to RCRA Regulatory Limits
NL Industries Site February, 1993

Element	Maximum Concentration Measured (mg/L)	RCRA Regulatory Limit (mg/L)
Arsenic	0.90	5.0
Barium	0.68	100.0
Cadmium	0.37	1.0
Chromium	0.096	5.0
Lead	460.0	5.0
Mercury	ND ⁽¹⁾	0.2
Selenium	ND ⁽²⁾	1.0
Silver	1.7	5.0

⁽¹⁾ Method detection limit of 0.002 mg/L (2) Method detection limit of 0.01 mg/L.

Table 7. Results of TOC Analysis of Soil and Sediment Samples, NL Industries Site, December, 1992.

Sample Location	TOC Concentration (mg/kg)
E17	> 30000 *
E12	> 30000 *
E29	> 30000 *
E16	> 30000 *
EE1A	> 30000 *
P1	3725
P2	2650
P3	7675
P4	4075
P5	6325
W18	> 30000 *
W20	> 30000 *
W12	> 30000 *
W2	> 30000 *
S4	> 30000 *
S2	> 30000 *
S13	> 30000 *
S11	> 30000 *

[•] Denotes that the subcontract laboratory reported that due to the nature of the sample, an appropriate quantity could not be used in the analysis to meet the upper detection limits.


TABLE 8. Results of Grain Size Analysis of Soil and Sediment⁽¹⁾
NL Industries Site
December, 1992

Sample	Sand					Total	Silt	Clay
Location	Very Coarse	Coarse	Medium	Fine	Very Fine	Sand		
E12	3.2	11.2	34.6	27.6	9.4	86.0	12.6	1.4
E16	2.8	10.4	33.1	27.3	8.1	81.7	16.0	2.3
E17	2.3	8.0	35.9	26.8	7.1	80.1	13.6	2.3
E29	2.7	10.1	.32.1	28.1	10.1	83.1	15.6	1.3
EE1A	8.7	11.6	13.1	13.2	6.3	52.9	41.3	1.5
P1	4.7	8.4	29.1	14.0	4.1	60.3	13.3	21.1
P2	6.6	14.7	24.1	13.0	12.3	70.7	10.1	15.8
Р3	1.8	9.8	35.5	28.1	8.4	83.6	12.9	2.7
P4	1.4	7.3	32.9	34.5	11.1	87.2	11.4	1.3
P5	3.4	6.0	19.7	19.6	10.9	59.6	16.5	17.2
S2	15.2	10.4	19.0	17.1	8.1	69.8	23.6	6.6
\$4	10.8	13.4	26.5	19.1	7.4	77.2	17.3	5.5
S11	10.0	10.1	27.3	26.1	10.9	84.4	14.0	1.6
S13	10.1	7.8	25.9	22.6	8.5	74.9	18.7	6.4
W2	7.0	9.6	29.4	25.7	8.2	79.9	17.1	2.9
W12	16.9	12.6	14.1	10.0	4.6	58.2	30.0	11.8
W18	1.4	5.3	16.1	12.2	7.4	42.4	31.5	21.8
W20	6.0	11.0	13.8	10.1	6.0	46.9	30.3	22.9

⁽¹⁾All values reported in percent of total sample.

Figures

NLI 002 0455



002

Appendix A

APPENDIX A
REAC ANALYTICAL REPORT
NL Industries Site
Final Report
February, 1993

ANALYTICAL REPORT

Prepared by Roy F. Weston, Inc.

National Lead Pedricktown, NJ

February 1, 1993

EPA Work Assignment No. 3-710
Weston Work Order No. 03347-033-001-4710-01
EPA Contract No. 68-03-3482

Submitted to M. Sprenger EPA-ERT

P. Bovitz Da

Task Leader

Vinor Kousal 1/2 V. Kansal /D

S. & A. Section Chief

W. S. Butterfield

Project Manager

Analysis by:

Prepared by: G. Karustis

Reviewed by: M. Barkley

Table of Contents

Introdu	ction		Page	1
Case Na	arrative		Page	1
Section	I			
	Analytical Procedure for Metals		Page	2
	Analytical Procedure for Metals in TCLP		Page	2
	Analytical Procedure for Total Organic Carbon		Page	2 2 2 3
	Analytical Procedure for Grain Size		Page	2
	Results of the Metals Analysis	Table 1.1	Page	3
	Results of the Metals in TCLP Analysis	Table 1.2	Page	
	Results of the Total Organic Carbon Analysis	Table 1.3	Page	
	Results of the Grain Size Analysis	Table 1.4	Page	
	Results of the Grain Size Analysis	Graph 1.1	Page	
Section	II			
	QA/QC for Metals		Page	26
	Results of the Duplicate Analysis	Table 2.1	Page	
	Results of the MS/MSD Analysis	Table 2.2	Page	
	QA/QC for Metals in TCLP		Page	29
	Results of the MS Analysis	Table 2.3	Page	
	QA/QC for Total Organic Carbon		Page	
	Results of the MS/MSD Analysis	Table 2.4	Page	
Section	III			
	Chain of Custody		Page	33
	Communications		Page	36
Append	lices			
	Appendix A Data for Metals and Metals in TCLP		Page	I31000
	Appendix B Data for Total Organic Carbon		Page	I35000
	Appendix C Data for Grain Size		Page	I36000
			~	

Appendices will be furnished on request

INTRODUCTION

REAC Laboratory, in response to Work Assignment # 3-710, provided analytical support for soil samples collected from the National Lead Site, Pedricktown, MD. This support included the analyses of environmental samples as summarized in the table below, QA/QC data review, and preparation of an analytical report. The analytical report includes a summary of the analytical methods, the results, QA/QC methods, and the QA/QC results.

The samples are summarized in the following table:

COC #*	Number of Samples	Sampling Date	Date Received	Matrix	Analysis	Laboratory
7532	9	12/8/92	12/16/92	Soil	Metals, TCLP Metals	ATEC**
7532	9	12/9/92	12/16/92	Soil	Metals, TCLP Metals	ATEC**
7533	9	12/8/92	12/16/92	Soil	Grain Size	ATEC**
7533	9	12/8/92	12/16/92	Soil	Total Organic Carbon	ATEC**
7534	9	12/9/92	12/16/92	Soil	Grain Size	ATEC**
7534	9	12/9/92	12/16/92	Soil	Total Organic Carbon	ATEC**

- * COC # denotes Chain of Custody number
- ** ATEC denotes ATEC Environmental Associates

CASE NARRATIVE

The subcontract laboratory stated that due to the peaty nature of the samples (a very high organic content), very small sample sizes were used in the total organic carbon analysis. This led to variability in the replicates. The organic content was so high that even with the small sample quantities, the range of the instrument was exceeded.

The subcontract laboratory reported the results of the metals analysis on an "as received" basis.

ANALYTICAL PROCEDURE FOR METALS

The subcontract laboratory determined the metals content of the samples using methods found in "Test Methods for Evaluating Solid Wastes", SW-846, 3rd Edition, November, 1986. The results are listed in Table 1.1.

ANALYTICAL PROCEDURE FOR METALS IN TCLP

The subcontract laboratory determined the metals content of the TCLP extracts using TCLP Extract Method 1311 found in the Federal Register 40 CFR, Part 261, Appendix II, 29 June 1990. The results are listed in Table 1.2.

ANALYTICAL PROCEDURE FOR TOTAL ORGANIC CARBON

The subcontract laboratory determined the total organic carbon content of the samples using Method 9060 found in "Test Methods for Evaluating Solid Wastes", SW-846, 3rd Edition, November, 1986. The results are listed in Table 1.3.

ANALYTICAL PROCEDURE FOR GRAIN SIZE

The subcontract laboratory determined the grain size distribution of the samples using ASTM Method D422. The results are listed in Table 1.4 and Graph 1.1.

Table 1.1 Results of the Metals Analysis
UA # 3-710 National Lead
(based on wet weight)

Sample ID	A18814	A18815	A18816	A18817	A18818	A18819	A18820	A18821	A18822	MDL
Location	E17	E12	E29	E16	W12	W2	EE1A	P1	P2	
% Solids	85	78	83	81	34	54	41	89	90	

	Conc	Conc	Conc	Conc	Conc	Conc	Conc	Conc	Conc	
Analyte	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)((mg/kg)
	*******	******					*******	*******	*******	******
Arsenic	5.4	4.4	5.0	6.8	19	7.0	5.2	5.4	3.0	2.0
Barium	6.8	18	9.0	9.2	26	32	14	23	13	2.0
Cadmium	ND	ND	ND	ND	ND	ND	MD	ND	MD	2.0
Chromium	7.5	8.4	7.8	8.8	7.3	4.8	38	9.5	16	2.0
Leed	420	380	270	440	780	100	380	170	96	2.0
Mercury	MD	ND	ND	ND	ND	ND	ND	ND	ND	1.0
Selenium	ND	MD	ND	MD	MD	ND	MD	ND	ND	2.0
Silver	ND	ND	ND	ND	ND	ND	ND	ND	ND	2.0
******		*****	*****	******			******			******

Sample ID	A18823	A18824	A18825	A18826	A18827	A18828	A18829	A18830	A18831	MOL
Location	P3	P4	P5	W18	W20	\$4	S2	\$13	S11	
% Solids	91	90	92	79	45	41	38	36	38	

	Conc									
Analyte	(mg/kg)	(mg/kg)(mg/kg)							
********	======		******	******	*******	******	*******	*****		
Arsenic	19	10	14	52	56	4.0	ND	50	30	2.0
Barium	27	12	30	89	47	24	10	31	15	2.0
Cadmium	2.2	3.3	20	23	4.6	2.1	MD	6.8	10	2.0
Chromium	9.8	6.7	14	26	20	7.6	5.0	6.9	4.9	2.0
Lead	3600	1000	16000	15000	5100	730	140	4900	2200	2.0
Mercury	ND	ND	MD	ND	ND	ND	ND	ND	ND	1.0
Selenium	ND	2.0								
Silver	ND	ND	ND	ND	NĐ	ND	ND	ND	ND	2.0

ND denotes Not Detected

MDL denotes Method Detection Limit

Table 1.2 Results of the TCLP Metals Analysis
HA # 3-710 National Lead

Sample ID	A18814	A18815	A18816	A18817	A18818	A18819	A18820	A18821	A18822	MOL
Location	E17	E12	E29	E16	W12	W2	EE1A	P1	P2	

	Conc	Conc	Conc	Conc	Conc	Conc	Conc	Conc	Conc	
Analyte	(mg/l)	(mg/l)	(mg/l)	(mg/l)	(mg/l)	(mg/l)	(mg/l)	(mg/l)	(mg/l)	(mg/l)
********	*******	EE28888	******	*******	******					******
Arsenic	ND	ND	ND	ND	ND	ND	ND	ND	ND	0.050
Barium	ND	0.12	ND	ND	0.11	1.0	ND	0.14	0.23	0.050
Cadmium	ND	ND	ND	ND	ND	ND	ND	ND	ND	0.010
Chromium	ND	ND	ND	NO	ND	ND	ND	ND	ND	0.050
Lead	0.55	0.72	0.42	0.57	0.90	0.24	0.45	0.48	2.9	0.050
Mercury	ND	ND	NO	MD	ND	ND	ND	ND	ND	0.002
Selenium	NO	ND	ND	ND	ND	NO	ND	NO	ND	0.010
Silver	MD	ND	ND	ND	ND	NO	ND	ND	ND	0.050
*******	*******	******			******	-			*****	******

Sample 1D A18823 A18824 A18825 A18826 A18827 A18828 A18829 A18830 A18831 MDL Location P3 P4 P5 W18 W20 S4 S2 S13 S11

	Conc									
Analyte	(mg/l)									
*******		******	*****	******	******	******		******	******	******
Arsenic	ND	0.85	0.50	0.050						
Barium	0.069	0.095	0.10	0.33	0.27	0.21	0.055	0.68	0.16	0.050
Cadmium	ND	0.049	0.29	0.37	0.026	ND	ND	0.074	0.066	0.010
Chromium	ND	MD	MD	0.096	ND	ND	ND	ND	ND	0.050
Lead	43	3.3	460	120	18	1.4	0.96	43	11	0.050
Mercury	ND	0.002								
Selenium	MD	ND	MD	0.010						
Silver	ND	MD	MD	ND	1.2	ND	ND	ND	1.7	0.050
		*****						******	******	

ND denotes Not Detected

MDL denotes Method Detection Limit

Table 1.3 Results of the Total Organic Carbon Analysis UA # 3-710 National Lead (based on wet weight)

Sample	ID Location	Conc	Range	Method
			of four	Detection
			Analyses	Limit
		(mg/kg)	(mg/kg)	(mg/kg)
	*********		**********	
Blank		ND		22
B18814	E17	> 30000 #		22
B18815	£12	> 30000 #		22
B18816	E29	> 30000 #		22
B18817	E16	> 30000 #		22
818818	W12	> 30000 #		22
818819	W2	> 30000 #		22
818820	EE1A	> 30000 #		22
818821	P1	3725 *	2200-4400	22
B18822	P2	2650 *	2400-3000	22
B18823	P3	7675 *	6800-8300	22
B18824	P4	4075 *	2900-5900	22
B18825	P5	6325 *	5600-7200	22
B18826	W18	> 30000 #		22
B18827	₩20	> 30000 #		22
B18828	54	> 30000 #		22
B18829	\$2	> 30000 #		22
B18830	s13	> 30000 #		22
B18831	\$13 \$11	> 30000 #		22

- * denotes that this value is the average of four replicates
- # denotes that the subcontract laboratory claimed that due to the nature of the sample an appropriate quantity could not be used in the analysis

Table 1.4 Results of the Grain Size Analysis WA # 3-710 National Lead

Sample ID Location	C18814 C	18815 C E12	18816 C E29	:18817 C E16	18818 C W12		18820 C	18821 C	18822 P2
Standard Sieve Size Sieve Size (u)					CPCT*		CPCT*		•. •.
0.5 in 0.375 in 4 4760 8 2380 10 2000 16 1190 30 590 40 420 50 297 80 177 100 149 200 74	99.8 97.7 85.7 69.1 49.8 28.7 23.0	99.1 96.8 85.6 70.0 51.0 30.0 23.4 14.0	99.6 97.3 87.2 73.2 55.1 34.0 27.0	99.3 97.2 86.8 72.5 53.7 32.5 26.4 18.3	99.0 83.1 70.5 63.9 56.4 49.0 46.4 41.8	99.0 93.0 83.4 71.4 53.9 34.3 28.2 20.0	96.7 91.3 79.7 72.2 64.6 56.8 53.4 47.1	99.6 97.6 94.7 93.6 90.0 81.6 69.5 52.5 41.3 38.5 34.4	97.9 97.7 95.5 92.3 91.1 85.7 71.0 58.3 46.9 38.2 33.9 21.6
Diameter (RM)	CPCT*	CPCT*	CPCT*	CPCT*	CPCT*	CPCT*	CPCT*	CPCT*	CPCT*
0.0356 0.0349 0.0347	********	12.5	******	*******		24.4	*****	222222	15.1
0.0346 0.0345 0.0341 0.0 <u>33</u> 8	13.3		16.3	13.3	38.4			28.6	
0.0329 0.0233 0.0226 0.0225 0.0222	8.3	7.5	10.3	7.3	18.5	5.1	50.5		15.1
0.0221 0.0217 0.0135 0.0134	3.3	2.4	2.3			2.9	28.2	26.7	
0.0133 0.0131 0.0126 0.0096		1.4		4.3	14.0	2.9	10.8	24.9	13.3
0.0095 0.0093 0.0090 0.0068	2.3	1.4	1.3	3.3	11.8	2.9	8.4	23.0	11.5
0.0067 0.0066 0.0064 0.0034	2.3	1,2	1.1	2.3	11.8	2.5	5.8	21.1	11.5
0.0033 0.0032 0.0014	0.7	0.7	0.2	1.6 0.7	11.4 4.8	0.4	7.8 4.2	18.9	11.1
0.0013								14.3	7.5


^{*} denotes Cumulative Percent Through

Table 1.4 Results of the Grain Size Analysis WA # 3-710 National Lead


Sample ID Location		C18823 (18824 C	18825 C P5	18826 C W18	18827 C W20	18828 C \$4	18829 C \$2	:18830 C \$13	18831 \$11
Sieve Size	Sieve Size (u)			CPCT*	••••	CPCT*		CPCT*	CPCT*	CPCT*
	0.5 in			99.4	96.4					
4	0.375 in 4760	99.4	100.0	98.3 95.7	95.9					
8	2380	99.2	99.8	93.3	95.7					
10	2000 1190	99.1	99.7	92.5	95.5 94.3	99.1 94.1	94.8	91.2	92.4	93.6
30	590	97.4 87.6	98.4 91.1	89.9 83.9	89.0	83.1	89.2 75.8	84.8 74.4	89.9 82.1	90.0 79.9
40	420	73.4	79.0	75.8	82.2	76.1	62.9	65.7	71.8	68.4
50 80	297 177	52.1	58.2 32.1	64.2	72.9	69.3	49.3	55.4 43.2	56.2	52.6 33.9
100	149	30.2 24.0	23.8	50.6 44.6	63.7 60.7	62.1 59.2	35.3 30.2	38.3	39.3 33.6	33.9 26.5
200	74	15.6	12.7	33.7	53.3	53.2	22.8	30.2	25.1	15.6
	Diameter									
	(mm)	CPCT*	CPCT*	CPCT*	CPCT*	CPCT*		CPCT*	CPCT*	CPCT*
	0.0368		4.7	*******	3823223	222222	******	******	******	*****
	0.0364		4.7							8.3
	0.0356						16.2			
	0.0354 0.0345			24.5				26.1	17.8	
	0.0332			24.3		47.3		20.		
	0.0325				42.8					
	0.0234 0.0232		2.7							6.4
	0.0228						12.3		12.1	0.7
	0.0224							18.3		
	0.0221 0.0218			20.8		32.6				
	0.0210				37.1	32.0				
	0.0136		0.7							
	0.0135 0.0133							9.5		3.6
	0.0132						10.4	- • •	10.2	
	0.0128			19.0	71 7	25.3				
	0.0124 0.0096		0.7		31.3					2.6
	0.0094		•				7.5	8.5	8.3	0
	0.0091			18.1		25.3				
	0.0089 0.0068		0.7		25.6					1.6
	0.0067		٠				5.5	6.6	6.4	
	0.0065			17.2	24 -	22.9				
	0.0064 0.0034	2.3	1.3		21.8		2.2	2.2		2.2
	0.0033					15.0			3.1	
	0.0032		.0.4	14.9	14.7		-0.4	4 7	4.7	-0.4
	0.0014 0.0013		-0.6	10.4	7.0	6.5	-0.6	1.3	1.3	-0.6
	*********		******				======	======	======	==±===

^{*} denotes Cumulative Percent Through


Graph 1.1 Results of the Grain Size Distribution for Sample C 18814 (E17) WA # 3-710 National Lead




Graph 1.1 Results of the Grain Size Distribution for Sample C 18815 (E12) WA # 3-710 National Lead



Graph 1.1 Results of the Grain Size Distribution for Sample C 18816 (E29) WA # 3-710 National Lead




Graph 1.1 Results of the Grain Size Distribution for Sample C 18817 (E16)
WA # 3-710 National Lead




Graph 1.1 Results of the Grain Size Distribution for Sample C 18818 (W12)


WA # 3-710 National Lead




Graph 1.1 Results of the Grain Size Distribution for Sample C 18819 (W2) WA # 3-710 National Lead




Graph 1.1 Results of the Grain Size Distribution for Sample C 18820 (EE1A) WA # 3-710 National Lead




Graph 1.1 Results of the Grain Size Distribution for Sample C 18821 (P1) WA # 3-710 National Lead




Graph 1.1 Results of the Grain Size Distribution for Sample C 18822 (P2) WA # 3-710 National Lead




Graph 1.1 Results of the Grain Size Distribution for Sample C 18823 (P3) WA # 3-710 National Lead




Graph 1.1 Results of the Grain Size Distribution for Sample C 18824 (P4) WA # 3-710 National Lead




Graph 1.1 Results of the Grain Size Distribution for Sample C 18825 (P5) WA # 3-710 National Lead




Graph 1.1 Results of the Grain Size Distribution for Sample C 18826 (W18) WA # 3-710 National Lead




Graph 1.1 Results of the Grain Size Distribution for Sample C 18827 (W20) WA # 3-710 National Lead



Graph 1.1 Results of the Grain Size Distribution for Sample C 18828 (S4) WA # 3-710 National Lead




Graph 1.1 Results of the Grain Size Distribution for Sample C 18829 (S2) WA # 3-710 National Lead




NLI 002 0483

Graph 1.1 Results of the Grain Size Distribution for Sample C 18830 (S13) WA # 3-710 National Lead



Graph 1.1 Results of the Grain Size Distribution for Sample C 18831 (S11)
WA # 3-710 National Lead



QA/QC FOR METALS

Results of the Duplicate Analysis

Samples A 18818 and A 18825 were chosen for duplicate analyses. The relative percent differences for sample A 18818 ranged from 1 to 15. Four relative percent differences were not calculated because the analyte was not detected in either analysis. The relative percent differences for sample A 18825 ranged from 12 to 44. Three relative percent differences were not calculated because the analyte was not detected in either analysis. The results of the duplicate analyses are listed in Table 2.1.

Results of the Matrix Spike/Matrix Spike Duplicate Analysis

Samples A 18818 and A 18825 were chosen for matrix spike/matrix spike duplicate (MS/MSD) analyses. The percent recoveries, listed in Table 2.2, ranged from 58 to 150. The relative percent difference (RPDs), also listed in Table 2.2 ranged from 0 (zero) to 11. Six percent recoveries and three relative percent differences were not calculated because the concentration of analyte in the sample was greater than the concentration spiked.

Table 2.1 Results of the Duplicate Analysis
WA # 3-710 National Lead

Sample ID A 18818

Analyte		Duplicate Analysis Conc (mg/kg)	Relative Percent Difference
Arsenic	19	22	15
Barium	26	27	4
Cadmium	NO	MD	NC
Chromium	7.3	7.4	1
Lead	780	680	14
Hercury	ND	ND	NC
Selenium	ND	ND	NC
Silver	MD	ND	NC

Sample ID A 18825

Analyte	Initial Analysis Conc (mg/kg)	Duplicate Analysis Conc (mg/kg)	Relative Percent Difference		
Arsenic	14	22	44		
Barium	30	39	26		
Cadmium	20	17	16		
Chromium	14	12	15		
Lead	16000	18000	12		
Mercury	ND	ND	NC		
Selenium	ND	ND	NC		
Silver	ND	ND	NC		

ND denotes Not Detected NC denotes Not Calculated

Table 2.2 Results of the MS/MSD Analysis for the Metals $\pm 4.4 \pm 3.710$ National Lead

Sample ID A 18818

	Sample	Spike	MS	Percent	MSD	Percent	RPD
	Value	Added	Value	Recovery	Value	Recovery	
	(ppm)	(ppm)	(ppm)	(MS)	(ppm)	(MSD)	
Analyte							
******	*******	*******	*******		******	*******	*******
Arsenic	9.375	50	63.00	107	64.00	109	2
Barium	26.42	200	227.3	100	226.9	100	0
Cadmium	0.6106	5.0	6.806	124	7.026	128	3
Chromium	7.328	20	28.29	105	28.81	107	2
Lead	780.9	50	795	NC	807.1	NC	NC
Mercury	0.24	2.5	2.355	85	2.24	84	1
Selenium	ND	5.0	5.60	112	5.50	110	2
Silver	ND	5.0	3.208	64	2.879	58	11
22232222	******	########	*****	*********	******	*******	

Sample ID A 18825

	Sample Value (ppm)	Spike Added (ppm)	MS Value (ppm)	Percent Recovery (MS)	MSD Value (ppm)	Percent Recovery (MSD)	RPD
Analyte	(pp.,)	(pp.m)	(ppm)	(110)	(plant)	(11.55)	
3222222		******		*******	*******	=======	222222
Arsenic	7.05	46.73	•	-	63.55	121	•
Arsenic	7.05	48.54	60.19	109	-	•	10
Barium	29.83	200	215.1	93	219.4	95	2
Cadmium	19.53	5.0	22.54	NC	23.94	NC	NC
Chromium	14.38	20	32.57	91	33.44	95	5
Lead	15930	50	12780	NC	16270	NC	NC
Mercury	0.038	2.380	2.743	114	•	•	•
Hercury	0.038	2.50			3.00	118	3
Selenium	MD	0.971	1.46	150			
Selenium	MD	0.935			1.40	150	0
Silver	0.3096	5.0	4.454	83	4.194	78	6

ND denotes Not Detected

NC denotes Not Calculated

QA/QC FOR METALS IN TCLP

Results of the Matrix Spike Analysis

Samples A 18818 and A 18825 were chosen for matrix spike (MS) analyses. The percent recoveries, listed in Table 2.3, ranged from 66 to 124.

The TCLP spikes were not run in duplicate and are bench spikes, not digestion spikes.

Table 2.3 Results of the MS Analysis for the TCLP WA # 3-710 National Lead

Sample ID A 18818

	Sample	Spike	MS	Percent
	Value	Added	Value	Recovery
	(mg/l)	(mg/l)	(mg/l)	(MS)
Analyte				
*******	*****	******	2722334	******
Arsenic	ND	0.050	0.051	102
Barium	0.108	1.0	1.251	114
Cadmium	ND	1.0	1.159	116
Chromium	ND	1.0	1.207	121
Lead	0.896	2.0	3.347	123
Hercury	ND	0.005	0.00543	109
Selenium	ND	0.050	0.042	84
Silver	ND	1.0	1.091	109
32222222			2222222	********

Sample ID A 18825

	Sample	Spike	MS	Percent
	Value	Added	Value	Recovery
	(mg/l)	(mg/l)	(mg/l)	(MS)
Analyte				
*******		******	******	*******
Arsenic	0.022	0.050	0.062	80
Barium	0.100	1.0	1.151	105
Cadmium	0.291	1.0	1.348	106
Chromium	ND	1.0	1.11	111
Lead	5.369	2.0	7.856	124
Mercury	NO	0.005	0.0058	115
Selenium	ND	0.050	0.033	66
Silver	ND	1.0	1.015	102
*******		******	******	*******

ND denotes Not Detected

RPD denotes Relative Percent Difference

QA/QC FOR TOTAL ORGANIC CARBON

Results of the Matrix Spike Analysis

Sample B 18822 was chosen for matrix spike/matrix spike duplicate analyses. The percent recoveries, listed in Table 2.4, ranged from 66 to 94. The relative percent differences, also listed in Table 2.4, were 5 and 23.

Table 2.4 Results of the MS/MSD Analysis for Total Organic Carbon WA # 3-710 National Lead

Sample	ID	В	18822
--------	----	---	-------

Sample Analysis (mg/kg)	•	MS Observed (mg/kg)	Percent Recovery	MSD Spike (mg/kg)	MSD Observed (mg/kg)	Percent Recovery	RPD
2680	3556	5344	75	3556	6036	94	23
2680	4444	5746	. 69	4324	5526	66	5

Roy F. Weston, Inc. Project Name: NATIONAL CEAN REAC, Edison, N.J. Project Number: 3347-033-00/- 47/0-0/ RFW Contact: P. BOVITE Phone: 000 Phone: (80) 632-9200 **EPA Contract 68-03-3482** ANALYSES HEQUESTED (6, TC PO. 15, 5 SAMPLE IDENTIFICATION TELP METALS AN HETALS **Date Collected** REAC # **Sampling Location** 12/8/12 AISCIH 16 or 64.55/48 EIT E/2 E 29 A 18816 EK6 2 A 18817 WIZF S A 18818 W2 A 18819 EEIA 5 A 18820 71 5 A 18821 5 118822 12/9/12 A 18823 O 118824 PY A 18825 S AR826 W18 W20 5 A18827 54 SD A 18828 AR829 56 52 A18130 5/3 5 b AB831 511 56 Special Instructions: # MS/45D FOR TCLIMETALS Magrix: FOR SUBCONTRACTING USE ONLY Sediment Potable Water AND AA HETALS FROM CHAIN OF DS - Drum Solids GW · Groundwater Water - TAKE TOLP AND AN METALS FROM S.AME MOZ DAK CUSTODY # **DL** - Drum Liquids SW · Surface Water 0 . Oil X - Other Słudge - REFER TO CONTRACT PARD 12MA2 ENCLOSE Homs/Resean Relinquished By Date Received By Hems/Reason Relinquished By Received By Time Date 3:30 12442

Roy r. Weston, Inc. REAC, Edison, N.J. **EPA Contract 68-03-3482** ORULAB WORN REQUEST

Project Number: <u>03347-033-00(- 47/0-0(</u>
Project Number: <u>03347-033-00(- 47/0-0(</u>
Phone: <u>(98) 432-9200</u>

NLT 002 0494

SAMPLE IDENTIFICATION

ANALYSES REQUESTED

REAC #	Sample No.	Sampling Location	Metrix	Date C	ollected	# of Bottles	Cont Prese	ainer/ rvetive	GRAINS 175	Toc	N	
	C18814	E17	S	12/1	1/12	1	32a64		メ			
	C18815	E12				1						
	C18816	E29										
	C18817	E N6		 		- -				<u> </u>		
	C 18818	W12		1 1		LL						
	C18819	W2		11						<u> </u>		
	C18820	EEIA	$\dashv \dashv$	1 1		├			 	<u> </u>	\perp	↓
	C18821	<u>Pl</u>		1-1					 	ļ	1	
	C18822			↓		_	<u> </u>		<u> </u>	 		d
	B18814	E/7		1 1		H	402 GLAS	15/4°C		<u> </u>		<u> </u>
	BNSIS	EIZ		 }-		Н-	 			 		
	B188/6	E 29	_	1		₩.				 		
	B 18817	E16	_ _ _	 		Н.	 			 	<u> </u>	
	B/8818	W12				∤ 			<u> </u>	1 - 1		
	B18819	W2		╁╼╼╂		├- ├-			ļ	 		
	B/8820	EEIA		╂		₩.	 		<u> </u>	 		++-
	B18821	P1 P2	- - -	╂		╁-}	1.	····	 	+ +	- 	
	B18822		<u> </u>	┼	<u> </u>	+	A		 	 		+
				+		 	 					
				Special		느 -			<u></u>	J		

DL - Drum Liquids

Water **Surface Water** Oil

FROM CHAIN OF

CUSTODY #

Nome/Resear	Relinquished By	Date	Received By	Date	Time	Hems/Reason	Relinquished By	Desk	Received By	Date	Time
ALC/ANHESS	SIRPA-	14strz	OlBen	13/6.52	3:30						
			0		—						Ī
		T									
				 							
					<u> </u>			1			
		1			†		·	1 1			

Roy F. Weston, Inc. REAC, Edison, N.J. EPA Contract 68-03-3482

Project Name: NATIONAL LEAD

Project Number: 03347-033-001-47/0-61

RFW Contact: P. BOVITE Phone: Q20 632-7200 CUSTOUT RECORD/LAB WORK REGUEST

NLI 002 0495

SAMPLE IDENTIFICATION

ANALYSES REQUESTED

REAC #	Sample No.	Sampling Location	Metrix	Date Collected	# of Bottles	Co	ntainer/ servative	GRAINSPE	Toc]
	C/8823	f3	5	12/9/92		3200 C	GLASS/4°C	X]
	C18824	[4	S		11_								
	C18825	P5	S							$\perp \lambda$]
	C1826	W18	Ş	 _	 						\		1
	C18827	W 20	S	L	$\sqcup \bot$			 	 		1		4
	C18828	<u>S4</u>	Sb			<u> </u>							4
	C18829	<u>sz</u>	20		 			 			\rightarrow	_	4
	C18830	\$13	50	 	├ -}-	1		 		_	-	 	4
	C/883/	SIL	50		$\sqcup \!\!\! \perp$	1	,	├ ──₩			$-\!$	v _	4
	B18823	<i>P</i> 3	5			402 60	45/4°C		X		\	8	1
	B18824	P4	5						 			1 8	4
	B/8825	P5	5	l				<u> </u>				1	4
	B18826	W/8	S	<u> </u>	- -			ļ				/ w	1~
	B18827	W20	S		- -			ļ				 \ 	د.
	B/8628	<u> </u>	50	 								 	2.22
	B18829	<u> </u>	50	 _ 					 			 	-
	B/8830	<u> </u>	50	ļ <u>ļ</u>	 	 	·	}	 			1	4
—	R 18831	\$11	56	<u> </u>	1	*	<u>, </u>	 	V			 	┨
						-						 	┦
				İ	<u> </u>	<u></u>							وجاد

SD - Sediment DL - Drum Liquids Surface Water

Water Oil

FOR SUBCONTRACTING USE ONLY FROM CHAIN OF CUSTODY #

	,	_	, ————————————————————————————————————	-			T				
Nems/Reason	Relinquished By	Date	Received By	Date	Time	items/Reason	Relinquished By	Dead	Received By	Date	Time
A4/ANA4515	ALCE	12/15/12	Quel Bry	12-16-52	3:30						
	ſ			T							
				<u> </u>							
								1 - 1			1
		 		 	 			 		 	



ROY F. WESTON, INC./REAC GSA RARITAN DEPOT 2890 WOODBRIDGE AVENUE BLDG. 209 ANNEX EDISON, NJ 08837-3679 908-632-9200 • FAX: 908-632-9205

ATEC Associates
5150 East 65th Street
Indianapolis, IN 46220-4871

December 14, 1992

Attn:

John Dwenger

Project # 3347-33-01-4710, National Lead

As per Weston REAC Purchase Order number 08-83468, dated 12-14-92, please analyze samples according to the following parameters:

Analysis/Method	Matrix	# of samples		
TCLP Metals/SW-846-1311 then analysis by SW-846-7000 Series	Soil	18		
Metals, TCLP list prior to extraction/SW-846-7000 Series	Soil	18	_	
Grain Size/ASTM D422 with hydrometer. See note below	Soil	18		
Total Organic Carbon/SW-846-9060	Soil	18	•••	
Data package as per attached Deliverables Requirements				

Samples are expected to arrive at your laboratory on December 15,1992. All applicable QA/QC(MS/MSD) analysis will be performed on our sample matrix. A preliminary data package including a signed copy of our Chain of Custody is due at REAC on December 30, 1992, with the complete data package due by January 15, 1993. For the grain size analysis, both % organic and % moisture should be included in to analysis.

Should any questions or problems arise concerning this project, please call John Johnson at (908) 603-8023. For any billing questions, please call Cindy Snyder at (908) 632-9200. Thank you.

Sincerely,

Misty Barkley

Analytical Projects Control Group Leader

Roy F. Weston, Inc. / REAC Project

MB:jj Attachments

CC.

R. Singhvi M. Sprenger

Central File

4710Con.Atec

V. Kansal

Subcontracting File

T. Scrittorale

C. Snyder

P. Bovitz

Misty Barkley



GSA RARITAN DEPOT 2890 WOODBRIDGE AVENUE **BLDG. 209 ANNEX** EDISON, NJ 08837-3679 908-632-9200 • FAX: 908-632-9205

ATEC Environmental Consultants 5150 East 65th Street Indianapolis, Indiana, 46220-4871

Attn: Mr. John D. Dwenger

Re: Project #3347-033-001-4710, National Lead Ref.#: 0193-03, ATEC Work Order Number 9212215 January 21, 1993

We are reviewing the data-for the analysis of As, Ba, Cd, Cr, Pb, Hg, Se and Ag. To complete our review. of this data we need the specific details of how each spike was performed: including the spike volume and concentration used along with the sample volume used.

Could you please have faxed to me at (908) 603-8041 the needed corrections by 1/22/93 and hard copy to follow immediately. Please telephone Chuck Shields at (908) 603-8042 if you have any questions.

Sincerely,

Misty Barkley

APC Group Leader

Roy F. Weston, Inc./REAC

CC:

C. Shields

P. Bovitz

Task Leader

Central File

#3-710

M. Spenger

Work Assignment Manager

Appendix B

APPENDIX B
U.S. EPA ERT/TAT XRF RESULTS
NL Industries Site
Final Report
February, 1993

00.

SPECTRACE 9000 (S/N Q023) LEAD X-RAY FLUORESCENCE DATA NATIONAL LEAD SITE PEDRICKSTOWN, NEW JERSEY 8-9 DECEMBER 1992 CONCENTRATIONS IN MG/KG (PPM)

SAMPLE	ID	DATE	LEAD		STANDARD DEVIATION
NLE12	8	DEC	372		120
NLE16	8	DEC	710		307
NLE17	8	DEC	1225		524
NLE1	8	DEC	192	J	35
NLE27	8	DEC	885		700
NLE29	8	DEC	523		148
NLE2	8	DEC	596		169
NLEE1A	8	DEC	3217		1445
NLP1	8	DEC	414		76
NLP2	8	DEC	229	J	83
NLP3	9	DEC	7035		1101
NLP4	. 9	DEC	3780		1104
NLP5	9	DEC	9998		2273
NLP6	9	DEC	6268		666
NLS11	9		5669		229
NLS13	9	DEC	17934		755
NLS2	9	DEC	272	J	20
NLS4	9	DEC	912		23
NLS8	9	DEC	915		157
NLW12	8	DEC	1407		22
NLW13	8	DEC	913		.24
NLW18	9	DEC	10475		74
NLW19	9	DEC	17193		61
NTMSO	9	DEC	19183		232
NLW22	9	DEC	13875		141
NLW2	8	DEC	734		15
NTM3	8	DEC	792		43

DETECTION LIMIT - 96
QUANTITATION LIMIT - 321
J - DENOTES LEAD CONCENTRATION BETWEEN
DETECTION AND QUANTITATION LIMITS

Draft Copy
Date 12/10/92

12-8-92	
0630 Arme office	
0700 REAL	
0730- leve REAC	
0900- Honve Site	
NL Industries	

ECAL	Expected	Obsived
Cd 109 Pb L	10.539	10.532
LB	12.613	12.622
	14,762	14.761
Emission Seule	77.16	12.110
Fe 55 5K	2.307.	2.3076
Emissi-fk.	5.89 6.89	5.8910
Am 741 /bis	10.539	10.552
LB	12.610	17,606
Emissi- 1K	59,56	59.533

Rescheck

Fe
(d 119 sané

Fé fenk

max but 3619

1/2 may 1810

1789-6.2689 2 7016 65093 2193-6.2842 795-6.5253

002 0501

11

	Measurement he		
," 	Cd 60 Sc		
·	Fe 30 Sic		
·: ·	Ar 30 sec		
	Sample	Pb_	
	GATS B	21,636	
	WHE 28	530	
	ZNL HW4	1643	
	NLE17-1		art simpling 1140
	NLEIT-2	1,062	
	NLE17-3	1,933	
	NLE12-1	541	
` <u>_</u>	NLE13-2	372	•
	NLE13-3	303	
	NLE 7-1	381	:
<u> </u>	NLEZ-Z	795	9
· ·	NLEX-3	612	;
·—	NLE16-1	839	
	NLE16-7	286	
<u>-</u>	NLE 16-3	1,005	
·—.	NLE 39-1	399	
·	NLE39-2	731	·
·	NLEJ9-3	439	3
	(QATS B RENAME Q	14(4) 46.29	12 Pacinistally
a—	SNL HE 28	53	3
,.—.·-	ENLHW-4	18.	
.;` ⁻	201275-B	30,9	
;—· ⁻	NLEI-1	31	
	NLEI-7		13
	NLE1-3		14
	1		
,			

			—_#
	Sample_	Pb	
	NLE27-1	1871	
	NLE 27-2	460	
	NLE 37-3	333	
	NLEE 1-A-1	2024	
	NLEE 1-A-2	+3 <i>18</i>	
	NLEEI-A-3	5250	
	WP - NIWZ	755	
	Niwz.Z	723	
70	NLWZ-3	722	;_
,	10EW 2 - 1	731	
	Niw3 Z	815	
	NLW3.3	829	
	WLWIZ-1	1397	
	NLW12-Z	. 1437	
	NLW12-3	1386	
<u> </u>	Niw13 - 1	941	
	NLWIZ3 -Z	552	
	NLU13-3	415	
	N'L PI-I	343	
	NI A-Z	Sic	
	NLA.3	37.5	
	NL 12-1	271	
	NLPZ Z	302	
7.6	NLF7-3	1/4	
414	QATSB	22101	
		21780	
·	GATS B	92047	
		519	
	NLHE28	483	
	NLHE 28	479	
	NLHW-Y	1629	
	Nrltw-4	(652	
	NCHW-4	1663	

NLI 002 0503

-		·	
∦	do who aled - data		
·	1900 - styped h	ide K Ind	el Coto
	29/		
	12.9.92		
	10-1-12		
	NL treligies	11. 1 1	
	(10 Net at	Idel willy	
	NL Fred dies C700 Met at C810 Hillord Sit	i trical o	~
	XLF.		
- - [Ecul	Expector	Cherred
	Cil 109 36 L-	10.539	10.530
		12.613	13.611
	<u>13</u>	14.767	14.767
	Ginnistian Park	<u> </u>	23.10
	Fe 55 SK.	2.301	3.30
	Emission Penk	5.89	5.697
├ ौ	Am 341 36 La	0539	10.518
	L'A	12.613	12,611
	Emmissie- Deak	5950	59.51
	•		
	REGCHIC		
J	Fe		
	Colog. Source		
	FE PLAK	77	<u> </u>
	max Count 34	73 6 5473)
- -	1/2 max 173	6.1	
- -			
	4 6.35 26 KEV	<u> </u>	420
	1588	1491	<u>, </u>
	6.2687 Kev	6.5	359. 9
	1843	173	9
 			, 54
_	054-6-25 - 0-29	OK.	6 25
			- 25
-			

Mic-surency time	
(il 60	
Te 30	T
in 30	
Sungle	FB & Surple
(QPTS-B	21,571
NLHE 28	456
CNL HW-4	1,767
NLP3-1	8542
NLP3->	6.618
NLP3-3	5,943
NLP4-1 #	5,227
N- P4-2	3,563
N-P4-3	3,550
NLP 5-1 #	13,057
NLp5-2	7,613
NLP5-3	9,334
NLPE-1	5,378
NLP6-3	6,447
NL96-3	6,980
NLW18-1 *	10,461
NLW18-3	10,513
NLW 18-3	10,392
NLW19-1	17,149
NLW 19-2	17,152
NLW 19-3	17,279
(QA75-B	20,874
NLHE-28	<i>555</i>
NLHE-28 LNLHW-4	3,274 19.028
NLWAUTI *	19.028
NL W 30 - 3	- 19,511 19,011
NLW30-3	19.011
	<u> </u>

Sindle 3 since willing	<u>P5</u>
NLW2D-1	14,076
NL in チケーヌ	13,778
NLN 70-3	13,772
WONLS 7-1 *	300
NLS2-2	753
. Nr23-3	259
NL54-1 =	931
NL54-2	880
VNLS4-3 Writing is NLS4-	926
Cn75 B	21482
G1153	21251
G1173B	21677
NIHEZS	453
NLHEZS	443
· NIHEZ 8	
ûp NIS8:1	716
NL 58- Z	930
NL38.3	1100
NL513.1 X	17,458
ML 513 L	17, 319
NL5133	18,997
NLHEZ8	513
NEHWY	1995
Nether	i073
Nettin	7151
(1) ON 511-1	5510
NLS11-Z	5505
N.51.3	5993

133

CHITSB 711176
WHEZS 476
NUMBER 1753
1700- left site 1900- arived REAC - K Findelicate

12-11-92 D800- Arrived office the Chem. My History Can I go to three shee led Silve rept Week W Reg Z THT

NCI 002 0207

Appendix C

LI 002 05

Appendix C

APPENDIX C
STATISTICAL OUTPUT
NL Industries Site
Final Report
February, 1993

Model: MODEL1

Dependent Variable: PB_AA

Analysis of Variance

		Sum of	Mean		
Source	DF	Squares	Square	F Value	Prob>f
Model	1 15	2535198.02	152535198.02	9.735	0.0066
Error	16 25	0697160.43	15668572.527		
C Total	17 40	3232358.44			
Root MSE	3958.	35478	R-square	0.3783	
Dep Mean	2872.	55556	Adj R-sq	0.3394	
c.v.	137.	79907			

Parameter Estimates

		Parameter	Standard	T for HO:	
Variable	DF	Estimate	Error	Parameter=0	Prob > T
INTERCEP	1	553.258559	1192.9069118	0.464	0.6490
PB XRF	1	0.496490	0.15912559	3.120	0.0066

	Dep Var	Predict	Std Err		Std Err	Student		Cook's
0bs	PB_AA	Value	Predict	Residual	Residual	Residual	-2-1-0 1 2	D
1	380.0	737.0	1157.136	-357.0	3785.447	-0.094	1 1	0.000
2	270.0	816.4	1142.260	-546.4	3789.962	-0.144	1 1	0.001
3	440.0	905.8	1125.979	-465.8	3794.831	-0.123	1 1	0.001
4	420.0	1161.5	1082.233	-741.5	3807.538	-0.195	i i	0.002
5	380.0	2152.0	961.153	-1772.0	3839.890	-0.461	i i	0.007
6	170.0	756.8	1153.382	-586.8	3786.592	-0.155	i i	0.001
7	96.0	667.5	1170.451	-571.5	3781.351	-0.151	1 1	0.001
8	3600.0	4048.5	1006.248	-448.5	3828.320	-0.117	i i	0.000
9	1000.0	2430.0	943.714	-1430.0	3844.213	-0.372	i i	0.004
10	16000.0	5518.2	1260.731	10481.8	3752.217	2.794	****	0.440
11	100.0	915.7	1124.200	-815.7	3795.359	-0.215	i i i	0.002
12	780.0	1248.3	1068.392	-468.3	3811.445	-0.123	i i	0.001
13	15000.0	5756.5	1313.318	9243.5	3734.136	2.475	****	0.379
14	5100.0	10075.9	2490.087	-4975.9	3077.018	-1.617	j ***j	0.856
15	140.0	687.3	1166.619	-547.3	3782.535	-0.145	i i.	0.001
16	730.0	1005.1	1108.476	-275.1	3799.981	-0.072	ii	0.000
17	2200.0	3368.4	946.429	-1168.4	3843.546	-0.304	i i	0.003
18	4900.0	9455.3	2306.874	-4555.3	3216.661	-1.416	**	0.516

 Sum of Residuals
 8.185452E-12

 Sum of Squared Residuals
 250697160.43

 Predicted Resid SS (Press)
 370669831.06

UNIVARIATE PROCEDURE

Variable=DIFF_PB

	Home	ents			Quantiles(Def=5)			Exti	emes	
N	- 18	Sum Wgts	18	100% Max	14080	99%	14080	Lowest	0bs	Highest	Obs
Mean	1798.833	Sum	32379	75% q3	2840	95%	14080	-6000(10)	2840(5)
Std Dev	4896.437	Variance	23975092	50% Med	445	90%	13030	-4520(13)	3440(8)
Skewness	1.419955	Kurtosis	2.807687	25% Q1	134	10%	-4520	-10(1)	3470(17)
USS	4.6582E8	CSS	4.0758E8	0% Min	-6000	5%	-6000	130(15)	13030(18)
CV .	272.2007	Std Mean	1154.101			1%	-6000	134(7)	14080(14)
T:Mean=0	1.558644	Prob> T	0.1375	Range	20080	,					
Sgn Rank	53.5	Prob> S	0.0182	93-91	2706						
Num ^= 0	18	• •		Mode	-6000						

CORRELATION ANALYSIS

5 'VAR' Variables: AS BA CD CR PB

Simple Statistics

Variable	N	Mean	Std Dev	Sum	Minimum	Maximum
AS	17	34.259372	43.267801	582.409324	3.333333	138.888889
BA	18	42.447900	32.595966	764.062207	8.000000	110.126582
CD	8	14.685769	10.651385	117.486155	2.417582	29.113924
CR	18	20.327315	20.341924	365.891674	7.44444	92.682927
PB	18	4437.902577	6338.347551	79882	106.666667	18987

Pearson Correlation Coefficients / Prob > |R| under No: Rho=0 / Number of Observations

	AS	BA	CD	CR	PB
AS	1.00000	0.77567	0.39231	0.20000	0.63093
	0.0	0.0003	0.3364	0.4415	0.0066
	17	17	8	17	17
ВА	0.77567	1.00000	0.40290	0.32392	0.66826
	0.0003	0.0	0.3223	0.1897	0.0024
	17	18	8	18	18
CD	0.39231	0.40290	1.00000	0.24034	0.74875
	0.3364	0.3223	0.0	0.5664	0.0326
	8	8	8	8	8
CR	0.20000	0.32392	0.24034	1.00000	0.16106
	0.4415	0.1897	0.5664	0.0	0.5232
	17	18	8	18	18
PB	0.63093	0.66826	0.74875	0.16106	1.00000
	0.0066	0.0024	0.0326	0.5232	0.0
	17	18	8	18	18

CORRELATION ANALYSIS

2 'VAR' Variables: PB

PB_X


Simple Statistics

Variable	N	Mean	Std Dev	Sum	Minimum	Maximum
PB	18	2872.555556	4870.272117	51706	96.000000	16000
PB_X	18	39.327222	109.094793	707.890000	0.240000	460.000000

Pearson Correlation Coefficients / Prob > |R| under Ho: Rho=0 / N = 18

	PB	PB_X	
P8	1.00000	0.84303 0.0001	wit weight
PB_X	0.84303	1.00000	
	0.0001	0.0	

National Lead



NATIONAL LEAD LINEAR MODEL

Model: MODEL1

Dependent Variable: PB_LEACH

Analysis of Variance

		Sum of	Mean		
Source	DF	Squares	Square	F Value	Prob>F
Model	1	97745.74349	97745.74349	14.954	0.0014
Error	16	104582.71347	6536.41959		
C Total	17	202328.45696	r r		
Root MSE	80	0.84813	R-square	0.4831	
Dep Mean	39	9.32722	Adj R-sq	0.4508	
c.v.	20	5.57802			

Parameter Estimates

		Parameter	Standard	T for HO:	
Variable	DF	Estimate	Error	Parameter=0	Prob > T
INTERCEP	1	-13.764423	23.48674257	-0.586	0.5660
PB_MG	1	0.119632	0.03093638	3.867	0.0014

NATIONAL LEAD LINEAR MODEL

	Dep Var	Predict	Std Err		Std Err	Student			Cook's
0bs	PB_LEACH	Value	Predict	Residual	Residual	Residual	-2-1-0	1 2	D
1	0.5500	-7.8532	22.627	8.4032	77.617	0.108	1	l I	0.000
2	0.7200	-7.9362	22.639	8.6562	77.614	0.112	1 .	1 1	0.001
3	0.4200	-9.8728	22.913	10.2928	77.533	0.133	ļ]]	0.001
4	0.5700	-7.2659	22.546	7.8359	77.641	0.101	1	1	0.000
5	0.9000	13.6806	20.177	-12.7806	78.290	-0.163	1		0.001
6	0.2400	-11.5490	23.157	11.7890	77.461	0.152	1		0.001
7	0.4500	-2.6766	21.934	3.1266	77.816	0.040	1	1 1	0.000
8	0.4800	-11.4793	23.146	11.9593	77.464	0.154	1	1 1	0.001
9	2.9000	-12.4883	23.295	15.3883	77.419	0.199	1	1 1	0.002
10	43.0000	33.5626	19.114	9.4374	78.556	0.120	1		0.000
11	3.3000	-0.4719	21.658	3.7719	77.893	0.048	1		0.000
12	460.0	194.3	44.373	265.7	67.583	3.932	1	*****	3.332
13	120.0	213.4	48.878	-93.3855	64.400	-1.450	**		0.606
14	18.0000	121.8	28.604	-103.8	75.619	-1.373	**	[· [0.135
15	1.4000	7.5 36 0	20.754	-6.1360	78.139	-0.079	1	1 1	0.000
16	0.9600	-9.3569	22.839	10.3169	77.555	0.133	1	1	0.001
17	43.0000	149.1	34.183	-106.1	73.266	-1.448	**		0.228
18	11.0000	55.4964	19.509	-44.4964	78.459	-0.567	*	1 1	0.010

Sum of Residuals -1.49214E-13 Sum of Squared Residuals 104582.7135 Predicted Resid SS (Press) 200719.7266

NATIONAL LEAD LOG LINEAR MODEL

Model: MODEL1

Dependent Variable: LOG_PB

Analysis of Variance

		Sum		Mean		
Source	DF	Squar	es	Square	F Value	Prob>F
Model	1	46.200	66	46.20066	83.579	0.0001
Error	16	8.844	44	0.55278		
C Total	17	55.045	10			
Root MSE	0	.74349	R-s	quare	0.8393	
Dep Mean	1	.76099	Adj	R-sq	0.8293	
c.v.	42	.22002				

Parameter Estimates

		Parameter	Standard	T for HO:	
Variable	DF	Estimate	Error	Parameter=0	Prob > T
INTERCEP	1	0.606735	0.21598715	2.809	0.0126
PB MG	1	0.002601	0.00028449	9.142	0.0001

NATIONAL LEAD LOG LINEAR MODEL

	Dep Var	Predict	Std Err		Std Err	Student			Cook's
0bs	LOG_PB	Value	Predict	Residual	Residual	Residual	-2-1-0	1 2	D
1	0.4383	0.7353	0.208	-0.2970	0.714	-0.416	1 1	- 1	0.007
2	0.5423	0.7334	0.208	-0.1911	0.714	-0.268	1 1	1	0.003
3	0.3507	0.6913	0.211	-0.3407	0.713	-0.478] }	J	0.010
4	0.4511	0.7480	0.207	-0.2969	0.714	-0.416	1 1	1	0.007
5	0.6419	1.2034	0.186	-0.5616	0.720	-0.780	j *i	j	0.020
6	0.2151	0.6549	0.213	-0.4398	0.712	-0.617	*	1	0.017
7	0.3716	0.8478	0.202	-0.4762	0.716	-0.665	*	1	0.018
8	0.3920	0.6564	0.213	-0.2644	0.712	-0.371	1 1	İ	0.006
9	1.3610	0.6345	0.214	0.7265	0.712	1.020	1 1	**	0.047
10	3.7842	1.6357	0.176	2.1485	0.722	2.974	i i	****	0.262
11	1.4586	0.8957	0.199	0.5629	0.716	0.786	i i	* İ	0.024
12	6.1334	5.1300	0.408	1.0034	0.621	1.614	ĺ	***	0.562
13	4.7958	5.5452	0.449	-0.7494	0.592	-1.265	**	İ	0.461
14	2.9444	3.5544	0.263	-0.6100	0.695	-0.877	1 *1	- 1	0.055
15	0.8755	1.0698	0.191	-0.1944	0.719	-0.270	İ	Ì	0.003
16	0.6729	0.7026	0.210	-0.0296	0.713	-0.042	1 i	į	0.000
17	3.7842	4.1468	0.314	-0.3627	0.674	-0.5 38	*	ĺ	0.032
18	2.4849	2.1125	0.179	0.3724	0.722	0.516	i i	* j	0.008

Sum of Residuals 1.110223E-15
Sum of Squared Residuals 8.8444
Predicted Resid SS (Press) 11.8360

085	LOC	AS	BA	CD	CR	PB	PERC_PB
1	E17	6.353	8.000		8.8235	494.12	2.2262
2	E12	5.641	23.077	•	10.7692	487.18	2.9558
3	E29	6.024	10.843		9.3976	325.30	2.5822
4	E16	8.395	11.358	•	10.8642	543.21	2.0986
5	W12	55.882	76.471	•	21.4706	2294.12	0.7846
6	W2	12.963	59.259	•	8.8889	185.19	2.5920
7	EE1A	12.683	34.146	•	92.6829	926.83	0.9711
8	P1	6.067	25.843	•	10.6742	191.01	5.0259
9	P2	3.333	14.444	•	17.7778	106.67	54.3750
10	P3	20.879	29.670	2.4176	10.7692	3956.04	21.7389
11	P4	11.111	13.333	3.6667	7.4444	1111.11	5.9400
12	P5	15.217	32.609	21.7391	15.2174	17391.30	52.9000
13	W18	65.823	110.127	29.1139	32.9114	18987.34	12.6400
14	W20	124.444	104.444	10.2222	44.4444	11333.33	3.1765
15	S4	9.756	58.537	5.1220	18.5366	1780.49	1.5726
16	S2	•	26.316	•	13.1579	368.42	5.2114
17	\$13	138.889	86.111	18.8889	19.1667	13611.11	6.3184
18	S11	78.947	39.474	26.3158	12.8947	5789.47	3.8000

National Lead

085	PERC_AS	PERC_BA	PERC_CD	PERC_CR	PERC_PB
1					2.2262
2		10.4000			2.9558
3	•	•	•	•	2.5822
4		•		•	2.0986
5	•	2.8769	•	•	0.7846
6	•	33.7500	•	•	2.5920
7		•	•	•	0.9711
8	•	10.8348	•	•	5.0259
9		31.8462			54.3750
10	•	4.6511			21.7389
11		14.2500	26.7273	٠.	5.9400
12	•	6.1333	26.6800	•	52.9000
13		5.9931	25.4174	5.83385	12.6400
14	•	5.1702	5.0870	•	3.1765
15		7.1750	•	•	1.5726
16	•	4.1800	:	•	5.2114
17	12.9600	15.7935	7.8353	•	6.3184
18	12.6667	8.1067	5.0160	•	3.8000